

The Large Quasar Reference Frame (LQRF) - an optical representation of the ICRS

A.H. Andrei^{1,2,3}, J. Souchay³, N. Zacharias⁴, R.L. Smart⁵, R. Vieira Martins^{1,2}, D.N. da Silva Neto^{2,6}, J.I.B. Camargo²,
M. Assafin², C. Barache³, S. Bouquillon³, J.L. Penna¹, and F. Taris³

¹ Observatório Nacional/MCT, R. Gal. José Cristino 77, CEP20921-400, RJ, Brasil
e-mail: oat1@on.br

² Observatório do Valongo/UFRJ, Ladeira Pedro Antônio 43, CEP20080-090, RJ, Brasil

³ Observatoire de Paris/SYRTE, 61 Avenue de l'Observatoire, F-75014 Paris

⁴ US Naval Observatory, 3450 Massachusetts Av. NW, Washington, DC 20392

⁵ INAF/Osservatorio Astronomico di Torino, Strada Osservatorio 20, 10025 Pino Torinese

⁶ Centro Universitário Estadual da Zona Oeste, Av. Manuel Caldeira de Alvarenga 1203, CEP23070-200, RJ, Brasil

Received September 15, 1996, accepted March 16, 1997

ABSTRACT

Context. The large number and all-sky distribution of quasars from different surveys, along with their presence in large, deep astrometric catalogs, enables the building of an optical materialization of the ICRS following its defining principles. Namely, kinematically non-rotating with respect to the ensemble of distant extragalactic objects; aligned with the mean equator and dynamical equinox of J2000; and realized by a list of adopted coordinates of extragalactic sources.

Aims. The LQRF was built with the care of avoiding incorrect matches of its constituents quasars, homogenizing the astrometry from the different catalogs and lists in which the constituent quasars are identified, and attaining global alignment with the ICRF, as well as fully consistent source positions

Methods. Starting from the updated and presumably complete LQAC list of QSOs, the initial optical positions of those quasars are found in the USNO B1.0 and GSC2.3 catalogs, and from the SDSS DR5. The initial positions are next placed onto UCAC2-based reference frames, following by an alignment with the ICRF, to which were added the most precise sources from the VLBA and VLA calibrator lists - when reliable optical counterparts exist. Finally, the LQRF axes are inspected through spherical harmonics, to define right ascension, declination and magnitude terms.

Results. The LQRF contains 100,165 quasars, well represented across the whole sky, with 10 arcmin being the average distance between adjacent elements. The global alignment with the ICRF is 1.5 mas, and the individual position accuracies are represented by a Poisson distribution peaking at 139 mas in right ascension and 130 mas in declination.

Conclusions. The LQRF contains J2000 referred equatorial coordinates, and is complemented by redshift and photometry information from the LQAC. It is designed to be an astrometric frame, but it is also the basis for the GAIA mission initial quasars' list, and can be used as a test bench for quasars' space distribution and luminosity function studies.

Key words. catalogs – reference systems – quasars – data analysis – astrometry

1. Introduction

The establishment of the ICRS as the celestial reference system (Arias et al., 1995, Feissel & Mignard, 1998) answers not only the desiderata in terms of the choice of the physical model and coherence, but also of practicality when confronted with the earlier dynamical model. This is especially true since the system is materialized by the direction of quasi-inertial grid points, which are given by distant, extragalactic sources, instead of nearby, moving stars. However, for a long time after the establishment of the ICRS, the faintness of the extragalactic sources prevented a precise determination of optical positions for a large number of them. The VLBI radio technique answered this need, by providing precise positions. But again, such precise positions could only be derived for a limited number of strong and stable sources, and the ones with richer observational history have come to define the ICRF (Ma et al. 1998). Yet even this small number of sources enabled the VLBI to make important contri-

butions in defining both geophysics and radio astronomy standards.

In contrast, concerning extragalactic sources on the optical domain, poor astrometry and small number of known sources made with which the standard representation of the ICRF were assigned to the stellar catalog Hipparcos (Perryman et al., 1997). Currently several extensions of the HCRF (Hipparcos Catalog Reference Frame), densify this representation, reaching fainter magnitudes.

The situation concerning the quantity of optically recognized bright quasars as well as concerning their sky distribution has witnessed significant changes in recent years (Véron-Cetty & Véron, 2006, Souchay et al., 2008). At the same time, the science's demand for accurate astrometry of these objects has also increased. Examples include micro and macro lensing, binaries, and space density counts, as well as the requirements of space astronomy missions (Andrei et al., 2008) and the very improvement of the stellar catalogs, that requires for a dense mesh of fiducial points (Fienga & Andrei, 2002, 2004).

Therefore, the conditions are put together to combine the large number of quasars optically recognized, with the large, precise optical catalogs in order to produce an optical materialization of the ICRS in terms of its first principles, namely kinematically non-rotating with respect to the ensemble of distant extragalactic objects, aligned with the mean equator and dynamical equinox of J2000, and realized by a list of adopted coordinates of extragalactic sources. This article presents one such realization, hereafter termed Large Quasar Reference Frame (LQRF).

In the following section the input data is presented and reviewed, whereas in its final subsection a schematic of the data stream is presented. The data stream is detailed in Sects. 3, 4 and 5, where the analytical expressions used to derive the catalog's intermediary steps are shown. In Sect. 3 a local astrometric solution is employed to place all quasar input catalogs onto the UCAC reference frame. In Sect. 4 the positions derived in this way are aligned with the ICRF frame. In Sect. 5 local inhomogeneities are tackled by spherical harmonics fitting and local average redressing. The LQRF catalog is presented in Sect. 6, and a summary and follow-up perspectives are presented in the last section.

2. Input data and data flow

The LQRF is built from gathering, comparing, and adjusting the positions of quasars and their stellar neighborhood, extracted from several catalogs and lists. A brief summary of their content is presented in the following subsections. The role of each one is also mentioned, for the sake of introducing the general flow of the data treatment. The final subsection outlines the data flow, leaving the details for the following main sections.

2.1. LQAC

The Large Quasar Astrometric catalog (Souhay et al., 2008), hereafter LQAC, compiles the detections of 113,653 quasars from the original entries of the eight largest lists, with the surplus extracted from the latest edition of the Véron & Véron catalog (Véron-Cetty & Véron, 2006) after an analysis on the reliability of the input contributors. The LQAC contains extensive information on the physical properties of each object. In particular, it contains, whenever available, the redshift, the magnitudes of the objects at 9 different optical bands (u , b , v , g , r , i , z , J , and K), and the radio fluxes at 5 different frequencies (1.4GHz, 2.3GHz, 5.0GHz, 8.4GHz, and 24GHz). The position of each object is given by the most precise equatorial coordinate directly available from the input lists. Therefore, the precision is quite variable, ranging from the sub-milli-arcsec of VLBI determinations to the level of a tenth of arcsec of the optical determinations, up to reach nearly one arcsec of single dish determinations. On the other hand, there are no double entries, either from misrecognition or from double radio spots. Though the positions unevenness prevent them from characterizing an astrometric reference frame, the LQAC positions and magnitudes, are adopted as starting point for finding the objects in the dense optical catalogs presented next. Figure 1 presents the LQAC sky distribution and sky density.

2.2. USNO B1.0

The U.S. Naval Observatory B1.0 catalog (Monet et al., 2003), hereafter B1.0, is the latest one of the series of all-sky ultra dense catalogs issued by the USNO. It brings two epochs positions and




Fig. 1. Sky distribution (equatorial coordinates, N up, E right) of the quasars found in the LQAC catalog. The highest density regions indicate the SDSS DR5 contribution. The scale represents the number of sources in regions of 9 sq° shown on the plots.




Fig. 2. Sky distribution (equatorial coordinates, N up, E right) of the quasars found in the USNO B1.0 catalogs. Again, the highest density regions indicate the SDSS DR5 contribution. The scale represents the number of sources in regions of 9 sq° shown on the plots. Notice that the scale is slightly different from that of Fig. 1 relative to the LQAC.

three colors magnitudes (Johnston B, R and I) for 1,042,618,261 objects, up to about $V=20$. It results from scans of 7,435 Schmidt plates taken for six sky surveys and calibrated using Tycho-2 stars (Hog et al., 2000). The nominal astrometric accuracy is 200 mas, but the individual objects position precision has a median value of 120 mas along each direction of measurement. Noticing that the quasars forming the LQRF seat on the B1.0 fainter end, where astrometric precision and accuracy degrades, the LQAC objects were searched for in the B1.0, and a neighborhood of stars, i.e. their position, proper motion, and magnitude, was also collected around the found objects. Figure 2 presents the sky distribution and sky density of the 83,980 quasars recognized in the USNO B1.0




Fig. 3. Sky distribution (equatorial coordinates, N up, E right) of the quasars found in the GSC2.3 catalog. As before, the highest density regions mark the SDSS DR5 contribution. The scale represents the number of sources in regions of 9 deg^2 shown on the plots. Notice that the scale is slightly different from that of Fig. 2 relative to the B1.0 catalog.

2.3. GSC2.3

The Second Generation Guide Star catalog latest version (Lasker et al., 2008), hereafter GSC2.3, is an all-sky catalog derived from the Digitized Sky Survey (Taff et al., 1990) that the Space Telescope Science Institute and the Osservatorio di Torino have created from the Palomar and UK Schmidt survey plates. It contains position, proper motion, and magnitude (Johnston B, R, and I) for 945,592,853 objects, and is expected to be complete to $R=20$. The astrometric total error is quoted as smaller than 300 mas, while the relative astrometric error is better than 200 mas. In fact the astrometric residuals against external reference catalogs only become larger than 150 mas above $R=18$ at the fainter end. The astrometric calibration was done using Tycho-2 stars. As seen, there are overwhelming similarities between the GSC2.3 and the B1.0 raw data, which however were measured and treated by distinct methods, enabling a most natural combination of the independent results issued from each of them. So, as before, the LQAC objects were searched for in the GSC2.3, and a neighborhood of stars (position, proper motion, and magnitude) was collected around the found objects. Figure 3 presents the sky distribution and sky density of the 93,943 quasars recognized in the GSC2.3.

2.4. SDSS DR5

The Sloan Digitized Sky Survey (SDSS) data release 5 (Adelman-McCarthy et al. 2007), hereafter DR5, represents the completion of the SDSS-I project. It covers $8,000 \text{ sq}^\circ$, on the north Galactic cap ($b < 30^\circ$) and on three stripes towards the south Galactic cap, reaching up to $i=20.2$, to a total of 215 million unique objects. The astrometry is preferably referred to the UCAC catalog, or else to the Tycho-2 catalog. The RMS residuals per coordinate are 45 mas and 75 mas for reductions against the UCAC (Zacharias et al., 2000) and the Tycho-2 (Hoeg et al., 1997), respectively, with additional 30 mas systematic errors in both cases (Pier et al., 2003). The internal astrometric precision is mainly limited by seeing effects, and it is quoted around 35

mas in each coordinate. A point to notice is that quasars are signaled in the DR5, to a total of 74,869 entries with quasi-stellar object spectral class. As seen from Fig. 1, the large number of DR5 quasars represents an important feature of the LQAC input.

2.5. UCAC2

The second USNO CCD Astrograph catalog (Zacharias et al., 2004), hereafter UCAC2, is the latest release of the ongoing UCAC project, designed to observe the entire sky for R magnitudes between about 7.5 and 16. The observed positional errors are about 20 mas for the stars in the 10 to 14 magnitude range, and about 70 mas at the limiting magnitude of $R=16$. The UCAC2 is a high density, highly accurate, astrometric catalog (positions and proper motions) of 48,330,571 stars covering the sky from -90 to $+40^\circ$ in declination and going up to $+52^\circ$ in some areas. The northern limit is a function of right ascension. The astrometry provided in the UCAC2 is on the Hipparcos system. Positions are given at the standard epoch of Julian date 2000.0. The UCAC2 supersedes the Tycho-2 catalog, having a stellar density that is over 20 times higher. It is used here as the choice catalog to place the quasar positions collected from the catalogs described in the above subsections onto the Hipparcos system. Nevertheless, since the UCAC2 coverage does not attain the entire sky, two additional stellar frames are complementary used and are presented below..

2.6. UCACN

What is hereafter termed UCACN is a cutout of the preliminary UCAC positions around the LQAC quasars in the northern part of the sky that is not reached by the UCAC2 coverage (Zacharias, 2007). Although, as discussed further on, the astrometric accuracy of the UCACN positions and its stellar density granted no degradation on the results obtained using it, it will be dropped in future revisions of this work as soon as of the forthcoming release of the all-sky catalog UCAC3. The nominal astrometric accuracy of the UCACN stars is similar to that of the UCAC2 ones, but no proper motion data were released for these stars. Throughout this paper when referring to the UCAC2 and UCACN ensemble the designation UCAC is used.

2.7. 2MASS

The Two Micron All-Sky Survey point source catalog (Cutri et al., 2003), hereafter 2MASS, derives from uniformly scanning the entire sky in three near-infrared bands to detect and characterize point sources brighter than about 1 mJy in each band, with signal-to-noise ratio greater than 10, using a pixel size of $2.0''$, each point in the survey having been imaged six times. The detectors worked to a 3σ limiting sensitivity of $J=17.1$, $H=16.4$, and $K=15.3$. The 2MASS contains the position of 470,992,970 sources, but no proper motions. The astrometry is referred to the Tycho-2 catalog and it is accurate to 70-80 mas over the magnitude range of $9 < K < 14 \text{ mag}$. Comparative studies have shown that the 2MASS positions agree with those from common UCAC2 stars to within 10 mas (Zacharias et al., 2003), and that the 2MASS is compliant to the ICRF in the range 100-120 mas. Taking advantage of its all-sky coverage, very high stellar density, and consistent astrometry, the 2MASS catalog is used here as an independent vehicle to place the LQAC quasars recognized in the large, deep catalogs on the Hipparcos system.

2.8. ICRF-Ext2

The International Celestial Reference Frame 2nd extension (Ma et al., 1998, Fey et al., 2004), hereafter ICRF-Ext2, is the present materialization of the International Celestial Reference System (ICRS) at radio frequencies. It represents the basic frame with respect to which the position of any object in the celestial sphere should be measured. In its primary contents, the ICRF consists of 212 sources, called "defining", whose positions are independent of the classical planes (equator, ecliptic) and reference points (equinox), but consistent with the previous realizations of the Celestial System as the FK5. With the help of a considerable amount of VLBI observations, the individual positions of the sources were found to be accurate to within roughly 0.25 mas, while the stability of the reference axes attains a remarkable 20 μ as (micro-arcsec) accuracy. In the ICRF initial version 608 sources complemented the defining 212 sources. The entrance of the ICRF-Ext1 and of the ICRF-Ext2 brought 109 new sources in the catalog, thus leading to a total number of 717 radio-sources. Notice that in the ICRF-Ext2 catalog, a very small sample of objects are not quasars: 10 of them are categorized as AGN (Active Galactic Nuclei) and 10 of them are categorized as BL LAC (BL Lacertae). Although the LQAC a priori exclusively considers quasars for its compilation, these particular objects were retained thanks to their astrometric accuracy. The ICRF-Ext2 is used here to match the VLBI positions directly to the derived optical positions. Such match serves to two distinct steps: (a) to redress the global frame orientation towards the ICRF, and (b) to redress local departures of the derived quasar catalog from the ICRF.

2.9. VCS6

The Very Long Baseline Array (VLBA) Calibrator Survey (Petrov et al., 2006), hereafter VCS6, consists of a catalog containing milli-arcsec accurate positions of 3,910 extragalactic radio sources, mainly quasars. These positions have been derived from astrometric analysis of dual-frequency 2.3 and 8.4 GHz VLBA observations, on the framework of the International Celestial Reference System (ICRS) as realized by the ICRF. Thus, to the level of precision and accuracy of the optical positions determined here, the VCS6 sources represent the ICRF. In order to strictly validate such representation, 524 VCS6 sources with formal inflated errors larger than 10 mas were trimmed off. The VCS6 sources not belonging to the ICRF-Ext2 were added to it, to obtain an enlarged radio frame, which is used for the two steps mentioned in the ICRF-Ext2 subsection, that is, the global and local redressing of the optical quasar positions towards the ICRF.

2.10. VLAC

The Very Large Array (VLA) Calibrator list (Claussen, 2006), hereafter VLAC, consists of 5,523 radio flux, structure code, position, and associated error entries for 1,860 extragalactic sources. It is foremostly indicated as a reference tool for observations at the VLA, but the radio positions are accurate and compliant with the ICRF. Then, likewise in the previous subsection, the VLAC positions were added to the ICRF-Ext2 core data set, after trimming off 297 of them with errors larger than 10 mas. The combination of the ICRF-Ext2, VCS6, and VLAC will be here termed as the enlarged radio frame (ERF) and it represents the ICRF to within an uncertainty of 10 mas. The order of priority for the radio lists to compound the enlarged radio frame




Fig. 4. Sky distribution (equatorial coordinates, N up, E right) of the enlarged radio frame quasars. The filled symbols mark the sources for which an optical counterpart is found in at least one of the deep, dense source catalogs (B1.0, GSC2.3, DR5). The enlarged radio frame is formed, in order of priority of choice, by the ICRF-Ext2 (718 sources), the VCS6 (2,684 sources), and the VLAC (59 sources).

was, ICRF-Ext2 (718 sources), VCS6 (2,684 sources), and finally VLAC (59 sources). The 3,461 sources now represent the ICRF on a better distributed all-sky basis (Fig. 4), and those for which the optical counterpart position is determined will actually define the best adherence of the LQRF to the ICRF.

2.11. Data flow

The various origins of the sources compiled in the LQAC imposes first of all to refer their positions to a single frame in the ICRS system. After that, the quasar frame so formed itself is aligned with the ICRF. Finally, the position of each source is calculated by weighted average. This data flow can be sketched as follows.

1. The LQAC entries are all admitted.
2. The LQAC quasars are searched for, by a main criterion of position coincidence, in the B1.0, GSC2.3, and SDSS DR5 high density catalogs. Each individual match is kept and treated separately.
3. Around each of the individual quasars thus matched a stellar neighborhood is extracted from the pertinent high density catalogs.
4. The stellar neighborhood contents from the high density catalogs are searched for in the stellar catalogs UCAC2, UCACN, and 2MASS, as before by a main criterion of position coincidence.
5. Within each neighborhood, the stars common to the high density catalog and to the stellar catalog have their positions projected onto the tangential plane centered at the quasar. A local transformation function from the high density catalog to the stellar catalog is calculated, mimicking the complete plate solution polynomials.
6. At this point, for each quasar, typically, a number of positions are determined (e.g., from more than one high density

catalog, and local solutions from the two stellar catalogs). Each of these families of positions will still be handled separately.

7. Within each family, the quasars for which there is a position in the enlarged radio frame (that is, the combination of the ICRF, VCS6, and VLAC catalogs) are found, since they are so flagged in the LQAC, what as before follows mainly a criterion of positional coincidence.
8. The subsets so defined are used to calculate the global rotation and zero point corrections to obtain coincidence with the ICRF origin and axes. The global rotation and zero point corrections are next applied to the positions of all quasars of each given family.
9. Likewise the previous step, the optical and radio positions in subsets are combined using orthogonal functions, in right ascension, declination, and magnitude, in order to correct systematic local departures from the ICRF axes directions. And likewise the significant orthogonal functions are applied to the positions of all quasars of each given family.
10. Again the optical and radio positions in subsets are combined in small regions around each quasar to derive its correction for localized inhomogeneities.
11. For each position of each source in a given family, a total error is assigned that combines the formal error from the high density catalog where it appears, the error from the particular local correction, the error from the particular global rotation and bias, and the error from both the significant orthogonal function adjustments and the inhomogeneity corrections.
12. From the squared inverse of the total error, a weight is assigned to each determined position of a given quasar. The final position of each quasar is given as the weighted average of the determined positions. The error in the position is assigned according to an individualized function of the optical to radio disagreement.

The comparisons of the final positions against the existing corresponding positions in the enlarged frame is used to obtain the global properties of the LQRF.

3. The local astrometric solution

In this section the points 1 to 6 of Sect. 2.11 (data flow) are derived.

Starting from the LQAC input list of quasars, the first step to get their astrometric positions, in order to form a whole consistent reference frame, is to have them matched to the position of objects in the two large, deep all sky catalogs, the USNO B1.0 and the GSC2.3, as well as in the SDSS DR5. The basic matching criteria is of position agreement within 1 arcsec. This is equal to the poorest astrometric accuracy in the LQAC contributing surveys. It is also a safe threshold given the typical seeing and astrophysical plate resolution limits. Additionally a magnitude limit was imposed preventing the selection of objects brighter than $R=7$, nevertheless this threshold was not reached by any source. No proper motion limit was enforced because both the USNO B1.0 and the GSC2.3 warn against possible offsets of the proper motions zero points. A post check was made by determining local zero points of proper motion, given by the average motion. In comparison to the local zeros of proper motion, just 0.7% of the GSC2.3 selected objects show significant motion, while for the B1.0 selected objects just 0.1% of them show significant motion. To verify what would be the possibility of a chance hit, the input coordinates were varied by random values between 1 and 5 arcmin. In this way the same region of the true objects was being

Table 1. Initial optical to radio relationships^a.

Tracer	B1.0 to RadPos	GSC2.3 to RadPos	DR5 to RadPos
N	1962	2028	300
$\Delta\alpha\cos\delta$	-35.2 ± 4.7	$+32.9 \pm 5.5$	-4.5 ± 3.2
$\Delta\delta$	$+123.5 \pm 4.9$	$+39.5 \pm 4.7$	$+13.9 \pm 3.2$
A1	$+4.7 \pm 6.2$	-6.6 ± 6.7	-3.5 ± 4.7
A2	$+5.9 \pm 5.8$	$+13.1 \pm 6.3$	-19.4 ± 5.1
A3	$+42.1 \pm 5.7$	-29.0 ± 6.1	$+6.6 \pm 4.1$
A4N	$+123.0 \pm 5.3$	20.9 ± 5.7	1.6 ± 4.5
A4S	$+128.0 \pm 11.9$	134.1 ± 12.5	

^a For the catalogs USNO B1.0, GSC2.3, and SDSS DR5 relatively to the collected sample of sources with precise radio interferometric positions (Extended Radio Frame - RadPos, either from ICRF-Ext2, VCS6 or VLAC), are presented: the average offsets in right ascension and declination, the orientation angles (A1, A2, A3) relative to the standard trihedron of equatorial coordinates, and the equatorial biases (A4N and A4S, north and south of -20°). All values are in milli-arcsec (mas) and the number of sources is given in the first line.

swept. Keeping the search radius to 1 arcsec, in only 0.2% of the cases was verified a hit on the false coordinates. Moreover, in only 0.1% of cases was a second object found significantly close to the adopted match. These satisfying proportions change very little when the search radius is enlarged to 2 arcsec. While this is an added proof of the overwhelming correctness of the matches, it at the same time indicates that there is little advantage in adopting a larger radius, and certainly no gain to the confidence level attached to the cross-identifications.

In all, 83,980 quasars are recognized in the B1.0, and 93,943 quasars are recognized in the GSC2.3. The sky distribution follows closely that shown in Fig. 1. From the figure it is clear the large contribution of the SDSS DR5. For the later, 74,825 quasars are retrieved using the SDSS query server. The positions from the three catalogs were added independently to the input list. The final input data contains 100,165 quasars. Just small fractions of the total appear in only one of the catalogs (1,405 only in USNO B1.0, 4,425 only in GSC2.3, and 3,276 only in SDSS DR5), while 61,532 quasars appear in all three catalogs.

These three catalogs that bring the input positions for the recognized quasars are originally placed on the ICRF J2000 reference frame by different astrometric pipelines based on the Tycho-2 catalog. Their global orientation and equatorial bias relative to the ICRF J2000 frame are shown in Table 1. Local deviations are also apparent in Figs. 5-7.

In order to homogenize all the original different positions, a local astrometric reduction was performed based on the catalogs UCAC and 2MASS. The local correction follows the standard tangent plane astrometric reduction, taking the quasar original position as the central point of the rigorous gnomonic projection (Assafin, Vieira Martins & Andrei, 1997). Through it the catalogs coordinates are adjusted to the reference frames represented by the stellar catalogs. Three assumptions are implicit to the method. That the stellar catalogs deliver an improved, and similar, representation of the ICRF, either relatively to the two Schmidt plate surveys, as well as relatively to the photometric one. That in a very restricted neighborhood the deviation be-




Fig. 5. USNO B1.0 right ascension and declination offsets distributions relative to the equatorial coordinates axes and the R magnitude. The offsets are given in the sense of catalog minus the radio interferometric positions from the ICRF-Ext2, or the VCS6, or the VLAC.

tween the deep surveys and the stellar catalogs, whichever one it is originated from, can be modeled by simple relationships. And that the removal of such deviations can significantly improve the accuracy of the quasar positions found in the large surveys. Prior investigations support these assumptions (da Silva Neto et al., 2005).

3.1. The polynomial degree of the solution

On the functional side, the initial question to be tackled is on the polynomial degree of the local astrometric correction representation. The UCAC is the least dense of the involved catalogs, and it presents a typical density of 10 stars per 5×5 arcmin. Translated to the Schmidt plates scale this means that a first degree solution with 10 reference stars can be applied over a physical area as small as 25mm^2 . According to the adopted a priori assumptions, and given the sky density of the stellar catalogs, the simplest models seem to be well suited. To verify such procedure, a trial local reduction was performed for 29 quasars randomly distributed across the celestial sphere, at every one hour of right ascension, off the galaxy plane. Around each of them, a field of UCAC2 stars was selected in boxes of side 10 arcmin, which contained at least 30 stars. For these 29 fields of quasars,

complete polynomial solutions were tested, with orders varying from 0 to 3^{rd} , and a straight averages solution. The quasar positions were taken from the USNO B1.0 and all positions were transported to the USNO B1.0 plate epoch, using the proper motions of either catalog. Table 2 summarizes the mean results.

In the first half of Table 2 the average local corrections and the corresponding standard deviation are shown. Though the values only very loosely represent the systematic offsets between the USNO B1.0 and the UCAC2, what is important to note is the similarity between all the rows. In the second half of the table, the average internal errors of each solution and their dispersion are shown. Again all rows behave quite alike. Therefore, to keep the smallest the region where the local solution is derived, and to maximize the ratio of the number of equations to the number of parameters to find, the complete first degree polynomial was adopted. This solution also produces the smallest set of standard deviations. The zero degree solutions fared just as well as the others, so there would be no particular advantage on taking an even smaller region, while the first degree polynomial allows for some modeling, which is useful face to the large numbers of sources that the local solution is going to be applied to. Two formulations of the first degree polynomial are concurrently used, the complete one (six independent parameters) - hereafter the six




Fig. 6. GSC2.3 right ascension and declination offsets distributions relative to the equatorial coordinates axes and the R magnitude. The offsets are given in the sense of catalog minus the radio interferometric positions from the ICRF-Ext2, or the VCS6, or the VLAC.

parameter model, and the so-called four constants one (in which the X and Y axes are constrained to have the same scale) - here-after the four parameter model. Furthermore, notice that with the six parameter model the error in the central, pseudo-tangential point, which is always the position of the quasar, was also found to be the smallest.

3.2. The stellar catalogs and the size of the local region

As discussed above, the UCAC2 offers the desiderata required for the local astrometric solution: high stellar density, accurate global representation of the HCRF and hence of the ICRF, range of magnitudes up to 16, and small zonal and local frame biases. The stellar density supports using a region as small as 10 arcmin across, which is always within the overlap zone of the Schmidt plate surveys contributing to the B1.0 and the GSC2.3.

However, the UCAC2 has incomplete coverage in the northern celestial hemisphere. Two remedies are used to face such hindrance, the use of preliminary positions of northern UCAC stars (here called UCACN), and the use of the 2MASS catalog. In the former case, the UCACN positions are by construction on the same system of the UCAC2, and at the same precision.

The 2MASS Point Source catalog positions are compliant with the ICRS via the Tycho-2 catalog and expected to be accurate to 70-80 mas over the magnitude range of $9 < K_s < 14$ mag. At the limit $K_s = 16$, the accuracy is not better than 200 mas. As the 2MASS does not provide proper motions, these values are good for the mean epoch of observation, 1999.

The pitfalls due to the magnitude gap between the 2MASS objects and the brighter Tycho-2 reference stars degrade the final astrometric accuracy. The internal precision of the 2MASS individual positions is better, at the level of 40-50 mas in the $9 < K_s < 14$ mag range. The comparison between 5.2 million stars common to the UCAC2 shows standard deviations of 70-80 mas in both coordinates, for the $9 < K_s < 14$ mag range. For the common stars around the quasars extracted to define the local solution, the difference average and standard deviation are $\overline{\Delta\alpha\cos\delta} = +3$ mas, $\sigma_{\Delta\alpha\cos\delta} = 99$ mas, and $\overline{\Delta\delta} = +11$ mas, $\sigma_{\Delta\delta} = 95$ mas, all in the sense 2MASS.minus UCAC2. Furthermore, from the limited sample of 700 quasars from the enlarged radio frame found in the 2MASS catalog, the analogous statistics are $\overline{\Delta\alpha\cos\delta} = +0$ mas, $\sigma_{\Delta\alpha\cos\delta} = 144$ mas, and $\overline{\Delta\delta} = -2$ mas, $\sigma_{\Delta\delta} = 138$ mas.




Fig. 7. SDSS DR5 right ascension and declination offsets distributions relative to the equatorial coordinates axes and the R magnitude. The offsets are given in the sense of catalog minus the radio interferometric positions from the ICRF-Ext2, or the VCS6, or the VLAC.

On the positive side, for the same region, the 2MASS catalog presents about 10 times the number of reference stars that does the UCAC catalog. In the small regions used for the local solution, it represents a coherent local frame, from which the zonal bias can be removed after later comparisons against the enlarged radio frame sources. It is well aligned with the UCAC and with the ICRF, and represents an all-sky frame. The local solution was computed independently using UCAC and 2MASS reference stars. Each of the results contribute, again independently, to the weighted average used to compute the final LQRF quasar coordinates.

3.3. The analytical expression for the local correction astrometric solution

On establishing the analytical form of the local astrometric solution, a final point to be considered concerns the usefulness of a magnitude dependent term, a feature often present in Schmidt plates based surveys (da Silva Neto et al., 2000). In the present case, the R magnitude represents the obvious choice, mainly because it is common to all the involved catalogs, and closer to

the central wavelengths of the 2MASS bands. At the same time, due to the quasars redshift and to the reddening for the stars, coupled with CCD top sensitivity and wider plate effective pass-band, the signal-to-noise ratio is usually higher towards the red. An inspection on the magnitude dependency of the catalog minus enlarged radio frame residuals reveals a small linear trend, larger for the $\Delta\delta$ offsets than for the $\Delta\alpha\cos\delta$ offsets. The same trend is verified for both B1.0 and GSC2,3 increasing with magnitude for declination, and decreasing with magnitude for right ascension. The amplitudes get only to 10 mas in the significant cases. In what concerns the DR5, the magnitude dependence is of much smaller amplitude and opposite tendency. The possibility of a magnitude equation is not ruled out in the presentation of either of the two deep catalogs. However, the causes and ensuing description would be out of the scope of this work. Here, an additional obstacle would be to bridge the magnitude gap between the reference stars and the much dimmer quasars. All accounted, the magnitude investigation is left to be investigated by the orthogonal functions in direct comparison with the enlarged radio frame positions.

Therefore, the enforced local solutions solution are:

Table 2. Comparison between the models for the local astrometric solution^a.

Solution	$\overline{\Delta\alpha\cos\delta}$	$\sigma_{\Delta\alpha\cos\delta}$	$\overline{\Delta\delta}$	$\sigma_{\Delta\delta}$
3 rd degree	+42	126	-25	140
2 nd degree	+43	127	-23	136
1 st degree	+61	110	-34	120
0 degree	+42	126	-34	119
offset	+56	118	-45	130

Solution	$\overline{\Sigma_X}$	σ_{Σ_X}	$\overline{\Sigma_Y}$	σ_{Σ_Y}
3 rd degree	137	63	127	64
2 nd degree	134	66	125	64
1 st degree	135	68	120	65
0 degree	133	68	117	64

^a In the top half of the table, the average corrections and their standard deviations are presented. In the bottom half, the average internal error in the solutions and their dispersion are presented. All values are in milli-arcsecs. The difference between the "0 degree" and the "offset" solutions is just that in the former residuals larger than 2.5σ are removed and the solution is recalculated.

$$\xi_{SCat} - X_{QCat} = a X_{QCat} + b Y_{QCat} + c \quad (1)$$

$$\eta_{SCat} - Y_{QCat} = d Y_{QCat} + e X_{QCat} + f \quad (2)$$

In Eqs (1) and (2), "SCat" is either the UCAC2 (or even its preliminary northernmost part) or the 2MASS, and "QCat" designates the B1.0, the GSC2.3 or the DR5. The relations above are fit to the smallest set of common "SCat" and "QCat" stellar objects in a neighborhood larger than 10 arcmin but smaller than a box of size of 30 arcmin, to a minimum of 6 stars. The solutions are calculated independently for right ascension and declination. Whenever that minimum number of stars is touched, either at the start or after elimination of stars, the four parameter model is used - in that case, obviously combining the two equatorial coordinates solutions. In fact, as presented in the next section, the four parameter model was run for all quasars and it delivers solutions in excellent coincidence with the six parameter solutions. The analytical form of the four parameter model is presented in Eqs. (3) and (4), where "SCat" and "QCat" keep their afore-defined meaning.

$$\xi_{SCat} - X_{QCat} = g X_{QCat} + h Y_{QCat} + i \quad (3)$$

$$\eta_{SCat} - Y_{QCat} = h Y_{QCat} - g X_{QCat} + j \quad (4)$$

3.4. Applying the local astrometric solution

As discussed in the previous sections, the initial task of producing the LQRF consists of carrying out a local astrometric reduc-

tion using the most precise and accurate stellar catalogs representing the HCRF, through the Tycho-2 reference frame, in order to locally redress the quasars' position input from the high density, deep catalogs..

There, however, the question of epochs comes up. The stellar catalogs have a mean epoch very close to J2000, being J1998.79 (± 2.78 y) for the UCAC2, and J2003.34 (± 0.24 y) for the UCACN, the UCAC2 preliminary northernmost part, and J1998.87 (± 0.08 y) for the 2MASS. Also, the DR5 mean epoch is J2002.96 (± 1.45 y), raising no questions. The dilemma on whether the local reduction ought to be performed in the epoch of the quasar input position or directly in J2000 is actually only meaningful for the cases when the B1.0 and the GSC2.3 are locally reduced by the UCAC2. For them it would not be unreasonable to speculate that a reduction made at the quasar epoch would diminish the scatter in the stellar positions, on the supposition that the stellar images themselves were at the quasar epoch (i.e., the epoch of the plate where the quasar is found) because of the smallness of the used field. The B1.0 mean epoch is J1977.12 (± 5.49 y). On the other hand, it is verified that the B1.0 positions are given as an average of plates reductions, in such a way to jeopardize the hypothesis of common epochs, given the range of magnitudes and to some extent the two different types of objects. For the GSC2.3, in contrast, the positions always refer to one single highest quality plate, whenever possible the red one. But, again, the GSC2.3 mean epoch is J1992.5 (± 3.72 y) and therefore there is a balance between reducing modestly the stellar positions scatter and the errors brought in by faulty or biased proper motions.

In order to actually verify which way would be preferred, the local corrections for the sources belonging to the enlarged radio frame were evaluated both at J2000 and the quasar position epoch. A second trial was made by considering only the sources from the enlarged radio frame belonging to the three input catalogs and that could be reduced by all the stellar frames. This subsample contains 240 sources but enables a direct comparison between all the possible ways of treatment. The outcomes show no net advantage in working at the quasar epoch. For the B1.0 catalog, the number of sources for which the local reduction converges drops; there is a marginal gain on the number of stars entering in the reduction but the number of actually used stars remains the same; there is no diminishing on the standard deviations from the solution corrections, or from the optical minus radio offsets; and the removal of the equatorial bias is less efficient (by 98 mas). The external check actually shows a small increase of the average detachment to the representation of the ICRF origin. For the GSC2.3 there is no important loss in working at the quasar epoch, but no gain either. The number of sources for which the local reduction converges remains the same, and so does the number of entry and used stars. The standard deviation in the solution does not change, and for the optical minus radio positions the offset scatter improves only marginally (by 4.5 mas). The equatorial bias is less efficiently removed (by 40 mas). Exactly the same results are obtained when inspecting the subsample of common sources. In view of these results it was decided to work only at J2000 for the local corrections. Thus all pairs of input quasar catalogs and stellar catalogs are treated identically.

A second point concerning the issue of epochs has already been touched in the previous discussion. It concerns whether the local reductions made with the UCACN, that is for the northernmost sources, and with the 2MASS would be significantly worse than those made with the UCAC2. As seen, both of them have a mean epoch close to J2000, where the local corrections

are performed. For the 2MASS catalog, which is not originally presented as an astrometric frame, the number of equations of condition is substantially increased for a same patch of the skies as defined for the UCAC2. To test the performance of the different catalogs, we use the DR5, which was shown in Sect. 3 to have intrinsically smaller scatter relative to the ICRF. As before, the enlarged radio frame and its subset of all common sources are employed, and they provide responses much alike. The UCAC2 based reductions produce the best results as expected. The right ascension and declination standard deviation in the offsets to the radio frame is at 47.5 mas. We remark that this is already a gain relative to the DR5 catalog offsets that are at 59 mas, as produced from Table 1 values. The standard deviation in the solution itself is at 33 mas. The UCACN based reductions perform much alike, being actually slightly better on declination and 30 mas worse on right ascension. The optical minus radio standard deviation is 14 mas worse than that of the UCAC2. The 2MASS based reductions fare somewhat worse than the UCACN ones, being larger by 18 mas in the solution standard deviation, and worse by 6.5 mas in the optical minus radio standard deviations. In this case, it is worth to point out that the average optical to radio position offsets are much similar for the UCAC2 and the 2MASS (differing by just 5 mas). It is also to be stressed that, even though not bringing a direct improvement in accuracy as the UCAC2 based ones do, the UCACN and the 2MASS local reductions enable to place the DR5 positions on the same frame with the B1.0 and GSC2.3 reduced positions, without impairing their intrinsic precision.

Table 3 summons up all the applied local corrections.

As can be seen from Table 3, the different quasar input catalogs (codes B, G, or S) require different local corrections, while the three stellar reference catalogs (codes U, N, or T) deliver such corrections at similar accuracy. The local correction outputs (codes C) agree when calculated either by the six parameter or by the four parameter models (codes 6 or 4). The local corrections are generally larger than the remaining optical minus radio offsets (codes Δ), which supports their usefulness. The optical minus radio mean offset in all cases is improved relatively to the one obtained using original quasar catalogs positions. As a consequence, the mean offsets become similarly small.

At this point, the three quasar input catalogs and the three stellar reference frames deliver concurrent, independent sets of positions. And whenever the 1st degree polynomial solution fails to converge, the four parameter solution results can be adopted equivalently. There are thus up to twelve possible local solutions calculated for each source. For 61% of the sources more than one solution was obtained, for 844 sources all twelve solutions went through, and for 12 sources no solution converged. Figures 8 to 11 display the local corrections for the most critical cases, namely the B1.0 and the GSC2.3, as given by the UCAC and 2MASS frames. Comparing with the Figs. 5-6, it is already clear that the local corrections bring the quasar's positions onto the ICRF. The comparison between these figures and the initial plots (Figs. 5-7) show that the local astrometric solution significantly redress the input positions from the large catalogs. Notice that the outcomes from the UCAC and 2MASS stellar reference frames bear much in common. Likewise, the six parameter and the four parameter models derive coincident corrections (thus, only those from the six parameter model are shown in the figures). It is also interesting to note that there is no discontinuity between the zones where either the UCAC2 or the UCACN stellar frames were used.

4. The global orientation towards the ICRS

In this section the points 7 and 8 of the Sect. 2.11 (data flow) are derived.

Once the local correction is obtained, the next move is to tie the preliminary frame just derived to the ICRF. As discussed in Sect. 1, the ICRF alone is too sparse, and in particular the number of its bright optical counterparts, to suffice for a detailed tying. Taking advantage of the fact that the long base radio interferometric observations always combine data for several sources in different sessions, a radio position net considerably larger than the ICRF was formed by adding the VLBA calibrators and the VLA calibrators. In the case of the VLA, only sources with the highest astrometric precision are included. The accurate, ICRF representing, radio interferometric positions of the enlarged radio sample sources (ERF) are then compared with their optical counterpart from the stellar based preliminary frame derived from the local corrections. From the ICRF 718 sources are collected, while from the VSC6 and the VLAC, 2,684 sources and 59 sources are collected respectively.

The enlarged radio frame so gathered thus contains 3,461 sources. They are adequately represented in all of the three contributing catalogs as detailed in Table 3 (column *Nrad*). The sky distribution was shown in Fig. 4. From the ERF sources, 2,263 have an optical counterpart in at least one of the input catalogs. The average minimum distance between the ERF sources is 187 arcmin, with a mode of 70 arcmin.

The ERF radio positions are then compared with their optical counterpart from the stellar based preliminary frame derived by the local corrections. Initially the comparison entails the derivation of global orientation corrections about the equatorial coordinates standard triad, plus a correction for a bias relatively to the equator. Following (Arias, Feissel & Lestrade, 1988), the canonical equations relating the right ascension and declination offsets to the rotations (A1, A2, A3) about the equatorial axes (namely, X on the equator pointing to the conventional origin, Y on the equator perpendicular to X, and Z pointing to the conventional pole) are given by

$$\Delta\alpha \cos\delta = A1 \sin\delta \cos\alpha + A2 \sin\delta \sin\alpha - A3 \cos\delta \quad (5)$$

$$\Delta\delta = -A1 \sin\alpha + A2 \cos\alpha + A4 \quad (6)$$

Equations (5) and (6) are solved independently as a measure to assess the robustness of the corrections. In this case, the direction cosines A1 and A2 agree when calculated either from $\Delta\alpha \cos\delta$ or $\Delta\delta$ within 1 σ in most cases. Figure 12 shows the good agreement on A1 and A2 as calculated from the right ascension residuals, from the declination residuals, or from the combined residuals. The equatorial bias (which is discussed in detail in the next section) can also be independently derived as simply $\Delta\delta = A4$. Also for A4, either the straight average or the adopted combined solution agree well.

For the sake of gathering the largest number of equations of condition, the final values are derived by a combined matrix including the three rotation corrections, plus the representation of the equatorial bias.


4.1. The equatorial bias

Former investigations (Assafin et al., 2001) pointed out marked differences in the equatorial bias concerning the USNO A2.0 cat-

Table 3. Mean results from the local astrometric corrections^a.

Solution	Nq	$\overline{Ns}$	$\overline{Cacos\delta}$	$\overline{C\delta}$	Nrad	$\overline{\Delta\alpha cos\delta}$	$\overline{\Delta\delta}$
BU6	47822	13.0	-23.5 (106.2)	-145.3 (104.2)	1320	-43.5 (177.0)	+12.2 (171.6)
BN6	3023	14.1	+ 4.2 (45.6)	-230.0 (81.7)	123	-70.5 (136.6)	- 9.9 (121.6)
BT6	83951	52.8	+ 3.3 (107.9)	-178.8 (111.2)	1897	-18.2 (172.9)	-12.0 (167.5)
BU4	43991	13.8	-21.8 (105.1)	-147.8 (103.6)	1273	-41.9 (176.6)	+11.0 (171.8)
BN4	3095	14.1	+ 4.7 (45.8)	-229.1 (80.3)	124	-68.8 (136.6)	-11.4 (121.9)
BT4	84012	52.7	+ 3.2 (107.7)	-178.9 (111.1)	1896	-18.6 (172.2)	-12.9 (167.0)
GU6	76671	22.7	-58.4 (109.7)	- 8.8 (91.4)	1633	-27.2 (178.3)	+33.7 (166.2)
GN6	4131	16.3	-56.3 (51.8)	- 5.4 (87.8)	150	+21.8 (148.8)	-11.9 (136.5)
GT6	93976	119.1	-55.9 (135.4)	-27.3 (119.2)	1984	- 5.1 (162.3)	+ 0.4 (157.3)
GU4	77110	22.6	-58.2 (109.9)	- 8.9 (91.4)	1634	-26.5 (178.2)	+33.0 (165.5)
GN4	4356	16.2	-56.6 (51.5)	+ 6.2 (86.3)	154	+20.1 (150.0)	-14.0 (133.3)
GT4	93978	119.1	-55.8 (135.4)	-27.4 (119.3)	1983	- 5.2 (161.8)	+ 0.3 (157.4)
SU6	59184	30.0	+ 6.5 (22.3)	+14.2 (26.2)	241	+ 5.7 (44.3)	+25.6 (49.9)
SN6	4035	16.0	- 7.6 (24.2)	-26.8 (48.9)	37	-15.0 (45.7)	- 1.8 (37.7)
ST6	74838	167.6	+ 5.3 (49.4)	+20.8 (55.5)	301	+ 9.8 (65.9)	+30.1 (63.1)
SU4	59125	30.0	+ 6.4 (21.8)	+14.1 (25.8)	240	+ 6.2 (43.3)	+24.1 (47.6)
SN4	4036	16.0	- 7.5 (23.9)	-26.5 (48.3)	37	-14.7 (45.4)	- 0.8 (37.0)
ST4	74838	167.6	+ 5.4 (49.3)	+19.9 (55.2)	301	+ 9.7 (65.9)	+29.0 (63.3)

^a Comparative results given by all types of local correction designed. The solutions nomenclature is formed by one letter designating the quasar position input catalog, followed by one letter designating the stellar reference frame, and by the codes 6 and 4 informing whether the algorithm is of the six or four parameter model. Nq is the number of quasars for which a given local correction was possible and succeeded. $\overline{Ns}$ is the average number of catalog stars contributing to the solution. $\overline{Cacos\delta}$ and $\overline{C\delta}$ columns bring the mean corrections on right ascension and declination, and the standard deviations (in brackets). The final two columns show the average optical to radio offset, preceded by the number of sources entering the average ($Nrad$). All angular values are in milli-arcsec. See the text for detailed explanations on the solutions and samples.

**Fig. 12.** Coincidence of the A1 and A2 direction cosines as obtained either from the $\Delta\alpha cos\delta$ residuals, from the $\Delta\delta$ residuals, or from both. The value adopted was from the solution including both the right ascension and declination residuals.

alog. The point of discontinuity seats around -20° of declination, which is the boundary of the north and south plate surveys that contributed to that catalog. Although such marked discontinuity should not befall neither the B1.0 nor the GSC23 catalogs, they mostly share the same constituent surveys with the USNO A2.0.

In order to analyze whether a discontinuity remains in the data, Eq. (6) was solved for the catalogs B1.0 and GSC23, as lo-

cally corrected by the UCAC2 and the 2MASS frames, adopting a single A4 unknown, two A4 unknowns dividing the $\Delta\delta$ offsets at -20° of declination, and eighteen A4 unknowns to fit the $\Delta\delta$ offsets within declination strips 10° wide. The plots in Fig. 13 show features that are more complex than could be described by one A4 standard offset. Nonetheless, there is a coarse positive plateau southwards of declination -20° , and values averaging near zero northwards. The standard deviation in the residuals from the 18 strips solutions is indeed no smaller than that of the -20° boundary, 2 A4 terms solution. This testifies of an interplay that can be more adequately handled in the next step where harmonic terms are introduced. On the other hand, the solution contemplating a single A4 term fares the worst by a few mas, and is certain to provide the least appropriate description.

The verified A4 declination dependency, prompted to assess whether a similar right ascension dependency is present. In this case, no such dependency exists.

4.2. Rotation angles and equatorial bias

The final values of the A1, A2, and A3 direction cosines for the global orientation towards the ICRF, as well as the equatorial bias, were found by adjusting the optical minus radio residual between the positions evaluated from the local astrometric solution and those given by the enlarged radio frame. The values are presented in Table 4.

For the B1.0 and GSC2.3 northernmost sources, for which the local correction was made using the UCACN, separate values were derived for the rotation angle and the (northern only) equatorial bias. In what concerns the DR5 northernmost sources, though the number of UCACN treated sources is small, the pre-




Fig. 8. Equatorial coordinates distribution of the local astrometric corrections derived by the six constants model using the UCAC stellar reference frame onto the USNO B1.0 quasar positions. The clear points correspond to the quasars for which there are ERF radio positions.

cisions are high and so a complete solution including one equatorial bias term was sufficient. At large, for the DR5 the A1 and A2 values, although significant and representing true corrections to be applied to the quasar equatorial coordinates, must be interpreted with caution in terms of what they would represent for the DR5 frame orientation, because of the concentration of the DR5 sources in the direction of the right ascension of 12h.

5. The removal of zonal bias

In this section the points 9 and 10 of the Sect. 2.11 (data flow) are derived.

5.1. Harmonic functions

At this point, the optical frame was homogenized by the local corrections, and was globally aligned with the ICRF. The final point is to sweep its axes to check for any remaining zonal modulations. This was done for the right ascension, declination, and R magnitude dimensions. In a sense this is equivalent to straighten

up the axes from waves of corrugation either trapped in the stellar catalogs themselves, or that were present in the initial deep catalogs and could not be completely removed by the local corrections.

The problem, the quantity of assessing points, the sky coverage, the dimensions of dependencies, and the relationship from the less precise, corrugated frame to the more precise, correcting one, are all the same to the case treated by Schwan (1988) when deriving zonal corrections to the FK5 construction. Thus, the same treatment by spherical harmonics (Brosche, 1966) is used here. The right ascension and declination corrections are expressed by the linear combination of the terms of a series, in which the $c_{pnml} - th$ element is given by

$$\Delta c = C_{c_{pnml}} H_p(R') L_n(\sin \delta) F_{ml}(\alpha) \quad (7)$$

where Δc is one or the other equatorial coordinates offsets, since independent spherical harmonic series are derived for each of them; $C_{c_{pnml}}$ is the $c_{prml} - th$ coefficient to be derived, for the c equatorial coordinate series; $H_p(R')$ is the Hermite polynomial of degree p , function of the R magnitude normalized by




Fig. 9. Equatorial coordinates distribution of the local astrometric corrections derived by the six constants model using the 2MASS stellar reference frame onto the USNO B1.0 quasar positions. The clear points correspond to the quasars for which there are ERF radio positions.

the mean and standard deviation for the quasars; $Ln(\sin\delta)$ is the Legendre polynomial of degree n , function of $\sin\delta$; and $Fml(\alpha)$ is the Fourier term of order m on α and sign l .

The offsets are produced by comparing the equatorial coordinates from the previous step (global rotation) with the VLBI positions from the ERF. That is, 2,263 pairs are available. Again, this is similar to the quantity available for the Schwan's (1988) application. Therefore terms up to very higher order can be derived. However, the analysis from the earlier steps advises to some caution in defining the actually significant order up which to halt. The distribution in right ascension of the offsets from the original deep catalogs hints for a 2π term and smaller ripples (cf. Figs. 5-7), while the analysis of global rotation coefficients has shown no α dependence. In contrast, the analysis of the equatorial bias displays a δ dependence, coherent in bins of 10° (Fig. 13). Magnitude dependences are believed to be possible within the large quasar catalogs. They could to some extent be introduced by the local solution itself, because of the magnitude difference between stars and quasars. The analysis made in Sect. 3.3 points out to such dependences but of feeble amplitude. All accounted for, and in order to have representative quantities of

ERF objects with which to compute the coefficients of each term, it was adopted to extend to 9^{th} order the Lagrange ($\sin\delta$) polynomials and the Fourier (α) functions. The Hermite (R magnitude) polynomials were extended to 2^{nd} order, but just to the 5^{th} order of the other two terms. Given the poor sky distribution, no declination terms were used for catalog combinations including the UCACN frame, nor right ascension terms for catalog combinations including the DR5 quasar list. All the above choices were made to retain only truly significant terms derived from the ERF comparison, such that they could be confidently applied to the full set of quasars.

Since by definition the spherical harmonics are orthogonally independent, their statistical significance can be tested individually. Each term was initially tested separately. Only those statistically significant at 3σ with coefficients not smaller than 1 mas were retained. Once more, this taxes heavier on the high order polynomials. Next, starting from the lower orders and up, the corrections brought by each term are removed from the equatorial offsets, and the following term is individually tested. The same criterion of 3σ significance and coefficient not smaller than 1 mas is applied to retain the terms. Actually, at this round the




Fig. 10. Equatorial coordinates distribution of the local astrometric corrections derived by the six parameter model using the UCAC stellar reference frame onto the GSC2.3 quasar positions. The clear points correspond to the quasars for which there are ERFA radio positions.

uppest order Hermite magnitude polynomials, when convolved with the higher Legendre (declination) and Fourier (right ascension) orders, are retained only if the coefficient is not smaller than 5 mas. The terms surviving these two rounds are added by a linear combination to separately adjust the right ascension and declination offsets by least squares. Table 5 reports the number of terms surviving the rounds. Notice the important decreasing rate from the first to the second round, which justifies the rounds scheme. On the other hand, there is no dropping from the second round (one by one regimen) to the third round (least squares fit of all validated terms together), what shows that they are truly statistically significant.

Table 6 reports the coefficients derived at the final adjustment for each family of quasar catalog and stellar frame. As expected, the coefficients do not vary from the second round to the final adjustment, indicating that no secondary harmonic survived. There is only one significant term including the DR5. It relates to the 2MASS local correction and depends solely on declination (3rd order). This reflects the high quality of the DR5 positions and indicates that the local correction did not induce further derail upon the input quasar catalogs. Concerning the

UCACN there is also only one significant term. It relates to the GSC2.3 quasar catalog and is of high right ascension order (8th). The lack of UCACN harmonic terms reflects the confinement of the sky region where it was applied. The positions originated from the B1.0 required 42 harmonic terms (31 for the right ascension offsets), and those from the GSC23 32 harmonic terms (20 for the right ascension offsets). Considering the stellar reference frames, the UCAC2 required 32 harmonic terms (25 for the right ascension offsets), while the 2MASS required 42 harmonic terms (28 for the right ascension offsets). In all cases, the right ascension offsets required more harmonic terms. In contrast, the $\ln(\sin\delta)$ dominated terms in general correspond to larger coefficients, which worked to correct the deviations detected at the A4 equatorial bias analysis. The magnitude terms appear only to the 1st order, in 18 B1.0 related terms and 10 GSC23 related terms. Thus, magnitude terms are presented as the catalogs authors cautioned, but they are less important than the zonal terms. Figure 14 shows the histograms of the harmonic terms coefficients and associated errors. The errors resemble well a Poisson distribution. The coefficients peak around -6 mas for the right ascension offsets, and around +6 mas for the declination offsets.




Fig. 11. Equatorial coordinates distribution of the local astrometric corrections derived by the six parameter model using the 2MASS stellar reference frame onto the GSC2.3 quasar positions. The clear points correspond to the quasars for which there are ERF radio positions.

5.2. Local inhomogeneities

Figures 15 and 16 (open symbols) show that after the correction by harmonic series, the optical minus radio residuals are equal to zero within the statistical significance of $3 \times$ standard error, that is without being attached to the scatter of the optical position determinations. This is shown for the pairs combining the B1.0 (Fig. 15) and GSC (Fig. 16) quasar input lists, and the UCAC2 (squares) and 2MASS (circles) local stellar frames, which are the largest and have the most homogenous sky coverage of the combinations used here.

Notwithstanding this, some clumping is seen, without a clear pattern over the sky, that can be dealt with further. The ERF sky density is such that the mean distance from any given quasar to the closest ERF VLBI position is 1.7° . And, on average, there are 10 ERF VLBI positions within a radius of 4.4° around any given quasar, the largest distance being on average 6.3° . Under such conditions for each of the combinations of quasar list and stellar reference frame, hence for every quasar a correction for the local inhomogeneities towards the VLBI position can be obtained from the average of 10 ERF quasars. Within the ERF quasars ra-

dus, objects are removed one by one if both their right ascension and declination offsets are larger than 2σ . On average, 1.3 ERF quasars are removed, what shows the robustness of the local inhomogeneities corrections. This procedure is similar to that of overlapping circles (Taff et al., 1990), but no overlap is actually required since the autocorrelation between adjacent corrections fades into a characteristic value at a distance of 20° . Figure 17 illustrates the weakness of the autocorrelations by taking each quasar in turn as a pole source, and plotting the standard deviation in the inhomogeneity correction within growing distance rings. For nearby sources, i.e., very small rings, the inhomogeneities are much the same, and the standard deviation is accordingly small. Rapidly though, before reaching the distance of 20° , the standard deviation already reaches the ceiling of scatter characteristic of each family of input catalog and stellar reference frame solutions.

By determining in this way the correction for the local inhomogeneities, it is seen in Figs. 15 and 16 (filled symbols) that the clumps are clearly minimized. After applying the corrections by harmonic functions the optical minus radio offsets are typically




Fig. 13. Declination dependency of the equatorial bias (A4). The plotted averages correspond to bins 10° wide in declination. The shaded codes indicate the number of ERF quasars within the bins.




Fig. 14. Distribution of the coefficients and associated errors of the harmonic functions fitting the $\Delta\alpha\cos\delta$ and the $\Delta\delta$ residuals. All significant terms (first round - see text) are counted here.

null within 1.3σ , what is reduced further to within 0.7σ after applying the complementary local inhomogeneities corrections.

6. The LQRF catalog

The LQRF catalog right ascension and declination equatorial coordinates are calculated from the values obtained in the final step of the data treatment (removal of zonal bias) by a weighted average. In the average, for each quasar, the solutions from the three input lists (B1.0, GSC23, and DR5) and the three stellar reference frames (UCAC2, UCACN, and 2MASS) are combined. The weights are the inverse of the square root sum of the internal and external errors at each of the correction steps plus the formal errors of each entry quasar input list. In order though to not carry too high or too low a weight for any given combination, for each of them separately the errors have been assigned by quartiles. It is worth noticing that solutions derived with different weighting schemes or even without any weighting agree within 10 mas. This indicates that the goal of homogenizing the different inputs was accomplished.

Since the final positions originate from the average of multiple combinations, attaching the final error of them to the squared sum of the contributing errors would in fact privilege the sources with fewer solutions obtained. Likewise, if attaching the error to

Table 4. Direction cosines relative to the ERF after the application of the local astrometric corrections^a.

Solution	N	A1	A2	A3	A4	A5
BU6	2616	-33.7 ± 6.3	-8.0 ± 6.1	50.8 ± 5.3	40.5 ± 10.7	2.6 ± 5.3
BN6	244	-5.5 ± 12.9	31.8 ± 14.1	123.0 ± 25.2	1.5 ± 12.8	
BT6	3764	-23.8 ± 5.1	1.5 ± 4.7	23.7 ± 4.6	27.5 ± 9.7	-21.0 ± 4.3
BU4	2522	-37.5 ± 6.3	-4.9 ± 6.2	48.6 ± 5.4	42.1 ± 10.7	0.0 ± 5.4
BN4	240	-0.4 ± 12.9	34.7 ± 14.1	118.9 ± 25.4	-0.3 ± 12.8	
BT4	3762	-24.6 ± 5.1	1.4 ± 4.7	24.3 ± 4.6	24.7 ± 9.6	-21.7 ± 4.3
GU6	3238	-7.0 ± 5.8	-3.1 ± 5.3	30.4 ± 4.7	100.8 ± 9.3	15.2 ± 4.8
GN6	300	-7.2 ± 13.2	16.3 ± 14.2	-43.4 ± 25.1	-4.7 ± 13.1	
GT6	3940	-4.0 ± 4.7	-4.9 ± 4.4	8.7 ± 4.3	30.2 ± 8.7	-6.1 ± 4.0
GU4	3240	-7.7 ± 5.8	-2.5 ± 5.3	29.4 ± 4.7	99.1 ± 9.3	14.7 ± 4.8
GN4	308	-7.1 ± 13.1	15.7 ± 14.0	-40.4 ± 24.7	-6.8 ± 12.9	
GT4	3938	-3.7 ± 4.7	-6.0 ± 4.4	8.7 ± 4.3	30.1 ± 8.7	-6.3 ± 4.0
SU6	480	-5.3 ± 4.7	-14.6 ± 4.8	-4.8 ± 3.7	16.6 ± 4.1	
SN6	74	-5.0 ± 11.6	14.8 ± 13.9	32.0 ± 17.8	9.8 ± 13.1	
ST6	600	-13.9 ± 5.5	-14.6 ± 6.0	-5.1 ± 4.9	20.1 ± 5.3	
SU4	478	-5.5 ± 4.5	-12.8 ± 4.6	-5.2 ± 3.5	16.3 ± 4.0	
SN4	74	-6.2 ± 11.4	13.1 ± 13.8	33.2 ± 17.6	9.5 ± 13.0	
ST4	600	-14.3 ± 5.5	-14.2 ± 6.0	-4.7 ± 4.9	19.2 ± 5.3	

^a Direction cosines A1, A2, and A3 from the comparison of the equatorial positions as found for each family of solutions and as given by the radio interferometry for the ERF objects. The equatorial bias A4 (north of declination -20°) and A5 (south of declination -20°), when applicable, are also presented. The solutions nomenclature is formed by one letter designating the quasar position input catalog, followed by one letter designating the stellar reference frame, and by the codes 6 and 4 informing whether the algorithm is of the six or four parameter models. N is the number of equations used to derive the direction cosines and equatorial biases. All angular values are in milli-arcsec.




Fig. 17. Fast fading of the autocorrelation of the local inhomogeneities correction. In the figure is plotted the standard deviation of the corrections in samples taken in rings of increasing radius. The upper lines correspond to the USNO B1.0 and GSC2.3 families of solutions, while the lower lines correspond the SDSS DR5 families of solutions.

the dispersion in the multiple contributing solutions. It is more realistic to derive the final error in the equatorial coordinates of each source from the external comparison of the LQRF position with the radio position. Since in the construction of the LQRF the three correction steps focused on the achievement of an homogeneous frame, this is done by taking the average of the LQRF minus radio positions for the 10 closest ERF sources

around each quasar. As seen before, this implies a typical radius of 4.4° around any LQRF quasar. Figure 18 presents the histograms of the errors in the coordinates. The distributions can be reckoned to Poisson distributions peaking at 139 mas in right ascension and at 130 mas in declination.

The LQRF contains 100,165 sources. On average, every source has a neighbor within 10 arcmin. Figure 19 shows the sky distribution counts within squares of 10°. Empty boxes are found only in the Galactic plane towards the southern hemisphere. Figure 20 presents a vector map of the offsets versus the VLBI positions, in bins of side 30° and at least 5 quasars. The mean offset is 32.7 mas and their distribution appears as random, with the largest offsets corresponding to bins of only 5 or 6 quasars.

Figures 21 to 24 show the optical minus radio coordinates offsets. The comparison with the equivalent Figs. 5-7 illustrates clearly the evenness of the ICRF representation produced by the LQRF positions.

The LQRF catalog is available through CDS access (in electronic form via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/VizieR?source=I/313>). An extract of its first page appears in Fig. 25. Its columns contain

1. the J2000 right ascension (h,m,s);
2. the J2000 declination (d,°,");
3. the external right ascension error ($\times \cos \delta$) (mas);
4. the external declination error (mas);
5. the number of radio interferometry neighbor positions used to evaluate the external errors;
6. the R magnitude, taken from the LQAC;
7. the redshift, taken from the LQAC (blank if not defined);

Table 5. Number of significant harmonic terms in the three rounds of testing^a.

Solution	R.A.			DEC.		
	1 st round	2 nd round	3 rd round	1 st round	2 nd round	3 rd round
BU6	21	6	6	2	1	1
BN6	0	0	0	0	0	0
BT6	17	9	9	17	5	5
BU4	23	7	7	0	0	0
BN4	0	0	0	0	0	0
BT4	17	9	9	17	5	5
GU6	15	6	6	8	3	3
GN6	0	0	0	2	1	1
GT6	9	4	4	5	2	2
GU4	15	6	6	8	2	2
GN4	0	0	0	2	1	1
GT4	9	4	4	5	2	2
SU6	0	0	0	0	0	0
SN6	0	0	0	0	0	0
ST6	1	1	1	0	0	0
SU4	0	0	0	0	0	0
SN4	0	0	0	0	0	0
ST4	1	1	1	0	0	0

^a Harmonic terms surviving in the three rounds of testing for the most significant ones. The three rounds are explained in the text, and the third round corresponds to the adopted harmonic fit. All procedure was done independently for the right ascension and declination residuals between the solutions and the ERF radio interferometry positions. The solutions nomenclature is formed by one letter designating the quasar position input catalog, followed by one letter designating the stellar reference frame, and by the codes 6 and 4 informing whether the algorithm is of the six or four parameter model.

8. the right ascension offsets to the radio interferometry position (blank if there is no radio position) ($\times \cos\delta$) (mas);
9. the declination offsets to the radio interferometry position (blank if there is no radio position) (mas);
10. the J2000 radio interferometry right ascension (h,m,s) (blank if there is no radio position);
11. the J2000 radio interferometry declination (h,m,s) (blank if there is no radio position).

7. Summary and perspectives

The LQRF is build to derive an optical representation of the ICRF, retaining the ICRS concept of being defined by extragalactic objects. Starting from the LQAC entries, quasars were identified in the USNO B1.0, GSC2.3, and SDSS DR5 catalogs. The positions there assigned were made homogeneous with respect to the HCRF by applying local corrections obtained in small neighborhoods around the quasars, relative to the UCAC and 2MASS catalogs. The global orientation was then adjusted to the ICRF and the zonal departures were removed. The ICRF was represented by the ERF containing accurate long base interferometry radio positions, collected from the ICRF-Ext2, the VCS6, and the VLACalib, and selected to precision better than 10 mas.

The final LQRF J2000 equatorial coordinates were derived by weighted averages of the input catalog positions, once locally, zonally and globally corrected as indicated. The errors assigned to the LQRF coordinates reflect the departure from the ERF.

The Large Quasar Reference Frame (LQRF) formed in this way contains 100,165 objects, of which 2,142 have accurate radio interferometric positions. The comparison between the LQRF positions and the ERF positions indicates that the overall

orientation towards the ICRF, as represented by the Euclidean direction cosines is $A1=+2.1 \pm 3.8$ mas $A2=-0.9 \pm 3.5$ mas $A3=-2.6 \pm 3.4$ mas with zero equatorial bias to the level of 2.9 mas. The average offsets to the ICRF are $\Delta\alpha\cos\delta=+2.7 \pm 2.9$ mas and $\Delta\delta=+0.3 \pm 2.9$ mas, with standard deviations of $\sigma_\alpha=134.3$ mas and $\sigma_\delta=131.1$ mas. The internal errors are well described by a Poisson representation, peaking at 139 mas in right ascension and at 130 mas in declination.. The LQRF is planned to be maintained and enlarged in the future, as new versions of the LQAC and other quasar input catalogs appear, as well as to incorporate newer versions of the UCAC2 stellar reference frame.

Acknowledgements. This work would have not existed without the effort and fundibg involved in the projects of the USNO B1.0 (US Naval Observatory), the GSC2.3 (Space Telescope Institute and Osservatório di Torino/IBAF), the Sloan Digitized Sky Survey (Astrophysical Research Consortium for the Participating Institutions), the ICRF (International Earth Rotation Service), the VLBA Calibrator Survey programs (Goddard Space Flight Center NASA), VLA (National Radio Astronomy Observatory), the UCAC project (U.S. Naval Observatory), and the 2MASS-The Two Micron All Sky Survey (NASA - NSF). The authors acknowledge their use.

A.H.A. thanks CNPq grant PQ-307126/2006-0. J.I.B.C. acknowledges grant 151392/2005-6/CNPq and grant E-26/100.229/2008/FAPERJ. M.A. acknowledges grant E-26/170.686/2004/FAPERJ and grants 306028/2005-0 and 478318/2007-3 (CNPq).

References

- Adelman-McCarthy, J.K., Ageros, M.A., Allam, S.S., Anderson, K.S. J., Anderson, S.F., Annis, J., Bahcall, N.A., Bailer-Jones, C.A.L., Baldry, I.K., Barentine, J.C., Beers, T.C., Belokurov, V., Berlind, A., Bernardi, M., Blanton, M.R., Bochanski, J.J., Boroski, W.N., Bramich, D.M., Brewington, H.J., Brinchmann, J., Brinkmann, J., Brunner, R.J., Budavri, T., Carey, L.N., Carliles, S., Carr, M.A., Castander, F. J., Connolly, A.J., Cool, R.J.,




Fig. 15. Averaged $\Delta \alpha \cos \delta$ and $\Delta \delta$ optical (from the USNO B1.0) minus radio (from the ERF) residuals in units of their standard deviations. Open symbols refer to the values corrected by the harmonic functions. Filled symbols refer to the values further redressed by the local inhomogeneities corrections. In both cases the averages are non significant below 3σ . The actual significance averages are 1.2σ after the harmonics treatment and 0.8σ after the complementary local inhomogeneities corrections.

Cunha, Ca.E., Csabai, I., Dalcanton, J.J., Doi, M., Eisenstein, D.J., Evans, M.L., Evans, N.W., Fan, X., Finkbeiner, D.P., Friedman, S.D., Frieman, J.A., Fukugita, M., Gillespie, B., Gilmore, G., Glazebrook, K., Gray, J., Grebel, E.K., Gunn, J.E., de Haas, E., Hall, P.B., Harvanek, M., Hawley, S.L., Hayes, J., Heckman, T.M., Hendry, J.S., Hennessy, G.S., Hindsley, R.B., Hirata, C.M., Hogan, C.J., Hogg, D.W., Holtzman, J.A., Ichikawa, S., Ichikawa, T., Ivezić, Z., Jester, S., Johnston, D.E., Jorgensen, A.M., Juric, M., Kauffmann, G., Kent, S.M., Kleinman, S.J., Knapp, G.R., Kniazev, A.Yu., Kron, R.G., Krzesinski, J., Kuropatkin, N., Lamb, D.Q., Lampeitl, H., Lee, B.C., Leger, R.F., Lima, M., Lin, H., Long, D.C., Loveday, J., Lupton, R.H., Mandelbaum, R., Margon, B., Martinez-Delgado, D., Matsubara, T., McGehee, P.M., McKay, T.A., Meiksin, A., Munn, J.A., Nakajima, R., Nash, T., Neilsen, E.H.Jr., Newberg, H.J., Nichol, R.C., Nieto-Santesteban, M., Nitta, A., Oyaizu, H., Okamura, S., Ostrik, J.P., Padmanabhan, Ni., Park, C., Peoples, J.Jr., Pier, J.R., Pope, A.C., Pourbaix, D., Quinn, T.R., Raddick, M.J., Re Fiorentin, P., Richards, G.T., Richmond, M.W., Rix, Hans-Walter, Rockosi, C.M., Schlegel, D.J., Schneider, D.P., Scranton, R., Seljak, U., Sheldon, E., Shimasaku, K., Silvestri, N.M., Smith, J.A., Smolcic, V., Snedden, S.A., Stebbins, A., Stoughton, C., Strauss, M.A., SubbaRao, M., Suto, Y., Szalay, A.S., Szapudi, I., Szokdy, P., Tegmark, M., Thakar, A.R., Tremonti, C.A., Tucker, D.L., Uomoto, A., Vanden Berk, D.E., Vandenberg, J., Vidrih, S., Vogeley, M.S., Voges, W., Vogt, N.P., Weinberg, D.H., West, A.A., White, Simon D.M., Wilhite, B., Yanny, B., Yocum, D.R., York, D.G., Zehavi, I., Zibetti, S., Zucker, D.B., 2007, *ApJS* 172, 634.

Andrei, A. H., Assafin, M., Barache, C., Bouquillon, S., Bourda, G., Camargo, J. I. B., le Campion, J.-F., Charlot, P., Gontier, A.-M., Lambert, S., Osrio, J. J. Pereira, da Silva Neto, D. N., Souchay, J., Vieira Martins, R., 2008, *Proc. IAU Symp.* 248, 260.

Arias E. F., Charlot P., Feissel M., Lestrade J.-F., 1995, *A&A*, 303, 604.

Arias, E.F., Feissel, M., Lestrade, J.-F., 1998, *A&A*, 199, 357.

Assafin, M., Vieira Martins, R., Andrei, A. H., 1997, *A&A*, 113, 1451.

Assafin, M., Andrei, A. H., Martins, R. Vieira, da Silva Neto, D. N., Camargo, J. I. B., Teixeira, R., Benevides-Soares, P., 2011, *ApJ*, 552, 380.

Brosche, P., 1966, *Veroff Astron Rechen-Inst*, no17.

Claussen, M., 2006, *VLA Calibrator Manual*.

Cutri, R.M., Skrutskie, M.F., van Dyk, S., Beichman, C.A., Carpenter, J.M., Chester, T., Cambresy, L., Evans, T., Fowler, J., Gizis, J., Howard, E., Huchra, J., Jarrett, T., Kopan, E.L., Kirkpatrick, J.D., Light, R.M., Marsh, K.A., McCallon, H., Schneider, S., Stiening, R., Sykes, M., Weinberg, M., Wheaton, W.A., Wheelock, S., Zacarias, N., 2003, *The IRSA 2MASS All-Sky Point Source Catalog*, NASA/IPAC Infrared Science Archive.

da Silva Neto, Dario N., Andrei, A. H., Vieira Martins, R., Assafin, M., 2000, *AJ*, 119, 1470.

da Silva Neto, D. N., Andrei, A. H., Assafin, M., Vieira Martins, R., 2005, *A&A*, 429, 739.

Fey, A.L., Ma, C., Arias, E.F., Charlot, P., Feissel-Vernier, M., Gontier, A.-M., Jacobs, C.S., Li, J., MacMillan, D.S., 2004, *AJ*, 127, 3587.

Fienga, A., Andrei, A.H., 2004, *A&A*, 420, 1163.

Fienga, A., Andrei, A.H., 2002, *A&A*, 393, 331.

Feissel M., Mignard F., 1998, *A&A*, 331, 33.




Fig. 16. Averaged $\Delta \alpha \cos \delta$ and $\Delta \delta$ optical (from the GSC2.3) minus radio (from the ERF) residuals in units of their standard deviations. Open symbols refer to the values corrected by the harmonic functions. Filled symbols refer to the values further redressed by the local inhomogeneities corrections. In both cases the averages are non significant below 3σ . The actual significance averages are 1.3σ after the harmonics treatment and 0.7σ after the complementary local inhomogeneities corrections..

- Hoeg, E., Bssgen, G., Bastian, U., Egret, D., Fabricius, C., Gromann, V., Halbwachs, J. L., Makarov, V. V., Perryman, M. A. C., Schwendiek, P., Wagner, K., Wicenc, A., 1997, *A&A*, 323, 57.
- Hoeg, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwendiek, P., Wicenc, A., 2000, *A&A* 363, 385.
- Lasker, B.M., Lattanzi, M.G., McLean, B.J., Garcia, J., Greene, G., Hanley, C., Hawkins, G., Laidler, V.G., Loomis, C., Meakes, M., Morrison, J., Rosenberg, A., Sturch, C.R., White, R.I., Wolfe, D., Bucciarelli, B., Carollo, D., Casalegno, R., Drimmel, R., Ferrari, A., Guglielmetti, F., Morbidelli, R., Pannunzio, R., Sarasso, M., Smart, R., Spagna, A., Volpicelli, A., Zacchei, A., Albretch, M., Benvenuti, P., Fourniol, N., Mignani, R., Pirenne, B., Favata, F., Gillet, F.C., Puxley, P., Simons, D., Hartley, M., Morgan, D.H., Read, M.A., Tritton, S.B., Djorgovski, G., Brucatto, R.J., Reid, I.N., 2008, *AJ*, 136, 735.
- Ma, C., Arias, E.F., Eubanks, T.M., Fey, A.L., Gontier, A.-M., Jacobs, C.S., Sovers, O.J., Archinal, B.A., Charlot, P., 1998, *AJ*, 116, 516.
- Monet, D.G., Levine, S.E., Canzian, B., Ables, H.D., Bird, A.R., Dahn, C.C., Guetter, H.H., Harris, H.C., Henden, A.A., Leggett, S.K., Levison, H.F., Luginbuhl, C.B., Martini, J., Monet, A.K.B., Munn, J.A., Pier, J.R., Rhodes, A.R., Riepe, B., Sell, S., Stone, R.C., Vrba, F.J., Walker, R.L., Westerhout, G., Brucatto, R.J., Reid, I.N., Schoening, W., Hartley, M., Read, M. A., Tritton, S. B. 2003, *AJ*, 125, 984
- Perryman, M. A. C., Lindgren, L., Kovalevsky, J., Hoeg, E., Bastian, U., Bernacca, P. L., Crz, M., Donati, F., Grenon, M., van Leeuwen, F., van der Marel, H., Mignard, F., Murray, C. A., Le Poole, R. S., Schrijver, H., Turon, C., Arenou, F., Froeschl, M., Petersen, C. S., 1997, *A&A* 323, 49.
- Petrov, L., Kovalev, Y.Y., Fomalont, E.B., Gordon, D., 2006, *AJ*, 131, 1872.
- Pier, J.R., Munn, J.A., Hindsley, R.B., Hennessy, G.S., Kent, S.M., Lupton, R.H., Ivezić, Z., 2003, *AJ*, 125, 1559.
- Souchay, J., Lambert, S. B., Andrei, A. H., Bouquillon, S., Barache, C., Le Poncin-Lafitte, C., 2008, *A&A*, 485, 299.
- Souchay, J., Andrei, A.H., Barache, C., Bouquillon, S., Suchet, D., Baudin, M., Gontier, A.-M., Lambert, S., Le Poncin-Lafitte, C., Taris, F., Arias F.E., 2008, *A&A*, 485, 299.
- Schwan, H., 1988, *A&A*, 198, 363.
- Taff, L. G., Lattanzi, M. G., Bucciarelli, B., Gilmozzi, R., McLean, B. J., Jenkner, H., Laidler, V. G., Lasker, B. M., Shara, M. M., Sturch, C. R., 1990, *AJ*, 353, 45.
- Taff, L. G.; Bucciarelli, B.; Lattanzi, M. G., 1990, *ApJ*, 361, 667.
- Véron-Cetty, M.-P., Véron, P., 2006, *A&A*, 455, 773.
- Zacharias, N., Urban, S. E., Zacharias, M. I., Hall, D. M., Wycoff, G. L., Rafferty, T. J., Germain, M. E., Holdenried, E. R., Pohlman, J. W., Gauss, F. S., Monet, D. G., Winter, L., 2000, *AJ*, 120, 2131.
- Zacharias, N., McCallon, H.L., Kopan, E., Cutri, R.M., 2003, *IAU GA 25, JD16*, Extending the Icrf Into the Infrared: 2MASS-UCAC Astrometry.
- Zacharias, N., Urban, S.E., Zacharias, M.I., Wycoff, G.L., Hall, D.M., Monet, D.G., Rafferty, T.J., 2004, *AJ*, 127, 3043.
- Zacharias, N., 2007, personal communication.

159.8 84.5 0 4 16.1276 +46 15 17.970

The figure consists of two vertically stacked histograms. Both histograms have a y-axis labeled 'Counts' ranging from 0 to 25,000 in increments of 5,000. The x-axis for both ranges from 0 to 600 mas, with major ticks every 100 mas. The top histogram is titled 'LQRF external R.A. errors (mas)' and shows a distribution peaking at approximately 25,000 counts for errors between 100 and 150 mas. The bottom histogram is titled 'LQRF external DEC. errors (mas)' and shows a similar distribution, also peaking at approximately 25,000 counts for errors between 100 and 150 mas. Both distributions are roughly Gaussian-like, centered around 100-150 mas.




Fig. 18. Distribution of the external errors assigned to the right ascension and declination J2000 positions of the LQRF sources. The distributions can be reckoned to Poisson distributions peaking at 139 mas on right ascension and at 130 mas on declination.




Fig. 20. Vectorial distribution of the systematic deviations (north up, east right) of the LQRF to the ICRF within bins of 10° . The average value is 32.7 mas, graphically represented by the off map arrow. All large values however befall to bins with 5 or 6 sources only.




Fig. 22. Distribution on right ascension of the $\Delta\delta$ LQRF minus ICRF residuals. There is no systematics above 4 mas and the median is 105.3 mas.




Fig. 21. Distribution on right ascension of the $\Delta\alpha\cos\delta$ LQRF minus ICRF residuals. There is no systematics above 4 mas and the median is 108.3 mas.




Fig. 23. Distribution on declination of the $\Delta\alpha\cos\delta$ LQRF minus ICRF residuals. There is no systematics above 4 mas and the median is 108.3 mas.




Fig. 24. Distribution on declination of the $\Delta\delta$ LQRF minus ICRF residuals. There is no systematics above 4 mas and the median is 105.3 mas.

Table 6. Values of the coefficients of the adjusted harmonic functions^a.

Solution	Offset	Hp(R')	Ln(sin δ)	Fml(α)	Coefficient
BU6	$\Delta\alpha\cos\delta$	0	0	1 -1	$+18.7 \pm 5.5$
BU6	$\Delta\alpha\cos\delta$	0	0	1 +1	$+41.7 \pm 5.2$
BU6	$\Delta\alpha\cos\delta$	0	4	0 0	-21.0 ± 6.0
BU6	$\Delta\alpha\cos\delta$	1	0	0 0	-5.5 ± 2.7
BU6	$\Delta\alpha\cos\delta$	1	3	0 0	$+13.1 \pm 2.9$
BU6	$\Delta\alpha\cos\delta$	1	4	1 -1	$+8.9 \pm 2.8$
BU6	$\Delta\delta$	0	8	3 +1	-20.2 ± 6.0
BN6	$\Delta\alpha\cos\delta$	-	-	-	-
BN6	$\Delta\delta$	-	-	-	-
BT6	$\Delta\alpha\cos\delta$	0	0	1 -1	$+16.6 \pm 4.7$
BT6	$\Delta\alpha\cos\delta$	0	0	1 +1	$+38.0 \pm 4.2$
BT6	$\Delta\alpha\cos\delta$	0	3	1 +1	$+12.3 \pm 4.4$
BT6	$\Delta\alpha\cos\delta$	0	4	0 0	-15.1 ± 4.6
BT6	$\Delta\alpha\cos\delta$	0	4	4 -1	-18.2 ± 4.5
BT6	$\Delta\alpha\cos\delta$	1	0	0 0	-8.7 ± 2.1
BT6	$\Delta\alpha\cos\delta$	1	3	0 0	$+7.5 \pm 2.1$
BT6	$\Delta\alpha\cos\delta$	1	5	0 0	-7.0 ± 2.1
BT6	$\Delta\alpha\cos\delta$	1	5	1 +1	$+7.2 \pm 2.2$
BT6	$\Delta\delta$	0	6	1 -1	$+14.9 \pm 4.8$
BT6	$\Delta\delta$	0	8	3 +1	-17.3 ± 4.5
BT6	$\Delta\delta$	0	9	6 +1	-14.2 ± 4.6
BT6	$\Delta\delta$	1	3	1 +1	$+6.9 \pm 2.0$
BT6	$\Delta\delta$	1	3	3 -1	$+7.2 \pm 2.1$
GU6	$\Delta\alpha\cos\delta$	0	0	1 +	$+18.0 \pm 4.7$
GU6	$\Delta\alpha\cos\delta$	0	1	0 0	$+64.2 \pm 9.0$
GU6	$\Delta\alpha\cos\delta$	0	3	0 0	-36.3 ± 8.5
GU6	$\Delta\alpha\cos\delta$	0	5	0 0	$+22.0 \pm 6.1$
GU6	$\Delta\alpha\cos\delta$	0	9	0 0	$+19.9 \pm 5.4$
GU6	$\Delta\alpha\cos\delta$	1	0	1 +	-7.1 ± 2.4
GU6	$\Delta\delta$	0	1	0 0	-23.0 ± 6.1
GU6	$\Delta\delta$	0	2	4 +1	-18.6 ± 5.7
GU6	$\Delta\delta$	1	0	0 0	$+7.3 \pm 2.3$
GN6	$\Delta\alpha\cos\delta$	-	-	-	-
GN6	$\Delta\delta$	0	0	9 -1	-33.9 ± 1.1
GT6	$\Delta\alpha\cos\delta$	0	0	1 +1	$+22.4 \pm 4.0$
GT6	$\Delta\alpha\cos\delta$	0	1	0 0	$+26.0 \pm 4.6$
GT6	$\Delta\alpha\cos\delta$	0	3	0 0	-21.0 ± 4.4
GT6	$\Delta\alpha\cos\delta$	1	1	1 -1	-7.9 ± 2.2
GT6	$\Delta\delta$	1	0	0 0	$+6.8 \pm 2.1$
GT6	$\Delta\delta$	1	2	4 +1	$+7.0 \pm 2.1$
DU6	$\Delta\alpha\cos\delta$	-	-	-	-
DU6	$\Delta\delta$	-	-	-	-
DN6	$\Delta\alpha\cos\delta$	-	-	-	-
DN6	$\Delta\delta$	-	-	-	-
DT6	$\Delta\alpha\cos\delta$	0	4	0 0	-14.8 ± 3.8
DT6	$\Delta\delta$	-	-	-	-

^a As the results from the six constants and four constants local astrometric solutions are quite alike, only the results from the six constants solutions are displayed. The code for the solutions are as in the previous tables and the coefficients are in milli-arcsec.